

4.061 Simultaneous and time-resolved measurement of alpha-pinene ozonolysis products in gaseous and aerosol phases for the determination of gas-aerosol partitioning.

Presenting Author:

Satoshi Inomata, National Institute for Environmental Studies, Tsukuba, Ibaraki, Japan, ino@nies.go.jp

Co-Authors:

Tanimoto Hiroshi, National Institute for Environmental Studies, Tsukuba, Ibaraki, Japan

Kei Sato, National Institute for Environmental Studies, Tsukuba, Ibaraki, Japan

Yuji Fujitani, National Institute for Environmental Studies, Tsukuba, Ibaraki, Japan

Yu Morino, National Institute for Environmental Studies, Tsukuba, Ibaraki, Japan

Kiyoshi Tanabe, National Institute for Environmental Studies, Tsukuba, Ibaraki, Japan

Phil Brown, Radboud University Nijmegen, Nijmegen, The Netherlands

Ryoji Yajima, Hokkaido University, Sapporo, Hokkaido, Japan

Yosuke Sakamoto, Kyoto University, Kyoto, Japan

Jun Hirokawa, Hokkaido University, Sapporo, Hokkaido, Japan

Abstract:

We have developed a real-time measurement system to identify semi-volatile organic compounds (SVOCs) in both the gaseous and the aerosol phases using chemical ionization-mass spectrometry in order to estimate the gas-aerosol partitioning of each SVOCs. In our system, SVOCs in both the gaseous and the aerosol phases are measured by the same technique, which enable us to determine the partitioning of each SVOC from the ratio of ion signals, without knowing the concentration of each SVOC. This ability is a strong point of this approach, because most chemical species in SOAs are thought to be multifunctional, and determining their concentrations seems to be impossible. We show a novel system equipped with dual-proton transfer reaction-mass spectrometers for “real-time” measurements of the partitioning of each SVOC produced in alpha-pinene ozonolysis between the gaseous and the aerosol phases. Time profiles of the SVOCs in both the gaseous and the aerosol phases, *i.e.* “real-time” SVOC partitioning, are also discussed.